

Do you struggle with bladder leaks when you:



SNEEZE



EXERCISE



LAUGH



COUGH



WALK



Don't settle for frustrating urinary symptoms or incontinence pads.

Ask your doctor today if Bulkamid is right for you.



To learn more about Bulkamid, visit www.bulkamid.com

References:

1. Chapple C, Dmochowski R. Particulate Versus Non-Particulate Bulking Agents in the Treatment of Stress Urinary Incontinence. Res Rep Urol. 2019; 11: 299–310.
2. Itkonen-Freitas AM et al. Tension Free Vaginal Tape Vs Polyacrylamide Hydrogel Injection For Primary Stress Urinary Incontinence: A Randomized Controlled Trial. J Urol. Vol 203, 372 – 378, 2020.
3. Bulkamid Patient Information Leaflet
4. Brosche et al. Seven-Year Efficacy and Safety Outcomes of Bulkamid for the Treatment of Stress Urinary Incontinence. NeuroUrol Urodyn. Vol 40(1):502-508, 2021
5. Dwyer L et al. "Voice Your Choice": A Study of Women's Choice of Surgery For Primary Stress Urinary Incontinence. International Urogynecology Journal. 31:769–777, 2020
6. Magon, N et al. Stress urinary incontinence: What, when, why, and then what? J Midlife Health. 2011 Jul-Dec; 2(2): 57–64.

BULKAMID[®]




Ready to say goodbye to bladder leaks?

Relief starts with **Bulkamid**[®]

Bulkamid is a minimally invasive and long-lasting treatment option for bladder leaks caused by stress urinary incontinence (SUI).

You're not alone!



1 in 3 women will suffer from SUI at some point in their lives.⁶

Important Safety Information:

Indications: Bulkamid[®] Hydrogel is intended to be used as a urethral bulking agent for the treatment of female urinary incontinence where the stress component is significant. Bulkamid[®] Hydrogel is intended for female patients, above the age of 18 years that have failed conservative treatments.

Contraindications: Bulkamid[®] Urethral Bulking System is contraindicated in patients suffering from acute cystitis or urethritis, have active Herpes Genitalis, or have damaged tissue in the urethra.

Warnings: Patients receiving treatment interfering with blood coagulation have an increased risk of haematoma or urethral bleeding.

Precautions: The Bulkamid Urethral Bulking System is only to be administered by a qualified, trained physician. If the patient has undergone major dental work or surgery, the Bulkamid Hydrogel should not be injected until the patient is fully recovered. Patients with acute or chronic infection in other sites of the body must be treated with caution. Only patients with well-controlled diabetes should be considered for Bulkamid Hydrogel injection. The procedure may cause urinary tract infections and scratches in urethra and bladder. Prophylactic antibiotic is recommended. Safety and effectiveness of device has not been established in patients: During pregnancy, delivery, or lactation; On immunosuppressive therapy or with autoimmune diseases; Under 18 years, Fragile urethral mucosal lining, Urethral hypermobility, Detrusor overactivity, Known polyuria, Unevaluated haematuria.

Side Effects: The following side effects may be associated with the use of the device system: General side effects normally associated with any surgical implantation procedure or local anaesthesia. Postoperatively, transient symptoms such as dysuria, stranguria, haematuria, urinary tract infection, and acute retention may occur; Scratching of the urethral mucosal may occur during the procedure; Long-term side effects such as non-acute retention, abscess formation, fibrosis (tissue hardening), de novo urgency, and necrosis are possible, but rare. Results and experiences may vary and are unique to each patient. No promise or guarantee is made about specific results or experiences. Talk to your doctor to find out if Bulkamid is right for you and to discuss the potential risks and benefits.

For a complete listing of indications, contraindications, warnings and precautions, go to www.bulkamid.com/isi
© 2021 Axonics, Inc. This material contains registered trademarks, trade names, and brand names of Axonics, Inc. All rights reserved. 110-0256-013rA 10/2021



Emergo Europe B.V. Prinsessegracht 20
2514 AP The Hague The Netherlands



Stay Connected.
#Axonics | #Bulkamid



Do you struggle with bladder leaks when you:



SNEEZE



EXERCISE



LAUGH



COUGH



WALK



Don't settle for frustrating urinary symptoms or incontinence pads.

Ask your doctor today if Bulkamid is right for you.



To learn more about Bulkamid, visit www.bulkamid.com

References:

1. Chapple C, Dmochowski R. Particulate Versus Non-Particulate Bulking Agents in the Treatment of Stress Urinary Incontinence. Res Rep Urol. 2019; 11: 299–310.
2. Itkonen-Freitas AM et al. Tension Free Vaginal Tape Vs Polyacrylamide Hydrogel Injection For Primary Stress Urinary Incontinence: A Randomized Controlled Trial. J Urol. Vol 203, 372 – 378, 2020.
3. Bulkamid Patient Information Leaflet
4. Brosche et al. Seven-Year Efficacy and Safety Outcomes of Bulkamid for the Treatment of Stress Urinary Incontinence. NeuroUrol Urodyn. Vol 40(1):502-508, 2021
5. Dwyer L et al. "Voice Your Choice": A Study of Women's Choice of Surgery For Primary Stress Urinary Incontinence. International Urogynecology Journal. 31:769–777, 2020
6. Magon, N et al. Stress urinary incontinence: What, when, why, and then what? J Midlife Health. 2011 Jul-Dec; 2(2): 57–64.

BULKAMID®




Ready to say goodbye to bladder leaks?

Relief starts with Bulkamid®

Bulkamid is a minimally invasive and long-lasting treatment option for bladder leaks caused by stress urinary incontinence (SUI).

You're not alone!



1 in 3 women will suffer from SUI at some point in their lives.⁶

Important Safety Information:

Indications: Bulkamid® Hydrogel is intended to be used as a urethral bulking agent for the treatment of female urinary incontinence where the stress component is significant. Bulkamid® Hydrogel is intended for female patients, above the age of 18 years that have failed conservative treatments.

Contraindications: Bulkamid® Urethral Bulking System is contraindicated in patients suffering from acute cystitis or urethritis, have active Herpes Genitalis, or have damaged tissue in the urethra.

Warnings: Patients receiving treatment interfering with blood coagulation have an increased risk of haematoma or urethral bleeding.

Precautions: The Bulkamid Urethral Bulking System is only to be administered by a qualified, trained physician. If the patient has undergone major dental work or surgery, the Bulkamid Hydrogel should not be injected until the patient is fully recovered. Patients with acute or chronic infection in other sites of the body must be treated with caution. Only patients with well-controlled diabetes should be considered for Bulkamid Hydrogel injection. The procedure may cause urinary tract infections and scratches in urethra and bladder. Prophylactic antibiotic is recommended. Safety and effectiveness of device has not been established in patients: During pregnancy, delivery, or lactation; On immunosuppressive therapy or with autoimmune diseases; Under 18 years, Fragile urethral mucosal lining, Urethral hypermobility, Detrusor overactivity, Known polyuria, Unevaluated haematuria.

Side Effects: The following side effects may be associated with the use of the device system: General side effects normally associated with any surgical implantation procedure or local anaesthesia. Postoperatively, transient symptoms such as dysuria, stranguria, haematuria, urinary tract infection, and acute retention may occur; Scratching of the urethral mucosal may occur during the procedure; Long-term side effects such as non-acute retention, abscess formation, fibrosis (tissue hardening), de novo urgency, and necrosis are possible, but rare. Results and experiences may vary and are unique to each patient. No promise or guarantee is made about specific results or experiences. Talk to your doctor to find out if Bulkamid is right for you and to discuss the potential risks and benefits.

For a complete listing of indications, contraindications, warnings and precautions, go to www.bulkamid.com/isi
© 2021 Axonics, Inc. This material contains registered trademarks, trade names, and brand names of Axonics, Inc. All rights reserved. 110-0256-013rA 10/2021



Emergo Europe B.V. Prinsessegracht 20
2514 AP The Hague The Netherlands



Stay Connected.
#Axonics | #Bulkamid



Do you struggle with bladder leaks when you:



SNEEZE



EXERCISE



LAUGH



COUGH



WALK

BULKAMID®



Don't settle for frustrating urinary symptoms or incontinence pads.

Ask your doctor today if Bulkamid is right for you.



To learn more about Bulkamid, visit www.bulkamid.com

References:

1. Chapple C, Dmochowski R. Particulate Versus Non-Particulate Bulking Agents in the Treatment of Stress Urinary Incontinence. Res Rep Urol. 2019; 11: 299–310.
2. Itkonen-Freitas AM et al. Tension Free Vaginal Tape Vs Polyacrylamide Hydrogel Injection For Primary Stress Urinary Incontinence: A Randomized Controlled Trial. J Urol. Vol 203, 372 – 378, 2020.
3. Bulkamid Patient Information Leaflet
4. Brosche et al. Seven-Year Efficacy and Safety Outcomes of Bulkamid for the Treatment of Stress Urinary Incontinence. NeuroUrol Urodyn. Vol 40(1):502-508, 2021
5. Dwyer L et al. "Voice Your Choice": A Study of Women's Choice of Surgery For Primary Stress Urinary Incontinence. International Urogynecology Journal. 31:769–777, 2020
6. Magon, N et al. Stress urinary incontinence: What, when, why, and then what? J Midlife Health. 2011 Jul-Dec; 2(2): 57–64.




Ready to say goodbye to bladder leaks?

Relief starts with Bulkamid®

Bulkamid is a minimally invasive and long-lasting treatment option for bladder leaks caused by stress urinary incontinence (SUI).

You're not alone!



1 in 3 women will suffer from SUI at some point in their lives.⁶

Important Safety Information:

Indications: Bulkamid® Hydrogel is intended to be used as a urethral bulking agent for the treatment of female urinary incontinence where the stress component is significant. Bulkamid® Hydrogel is intended for female patients, above the age of 18 years that have failed conservative treatments.

Contraindications: Bulkamid® Urethral Bulking System is contraindicated in patients suffering from acute cystitis or urethritis, have active Herpes Genitalis, or have damaged tissue in the urethra.

Warnings: Patients receiving treatment interfering with blood coagulation have an increased risk of haematoma or urethral bleeding.

Precautions: The Bulkamid Urethral Bulking System is only to be administered by a qualified, trained physician. If the patient has undergone major dental work or surgery, the Bulkamid Hydrogel should not be injected until the patient is fully recovered. Patients with acute or chronic infection in other sites of the body must be treated with caution. Only patients with well-controlled diabetes should be considered for Bulkamid Hydrogel injection. The procedure may cause urinary tract infections and scratches in urethra and bladder. Prophylactic antibiotic is recommended. Safety and effectiveness of device has not been established in patients: During pregnancy, delivery, or lactation; On immunosuppressive therapy or with autoimmune diseases; Under 18 years, Fragile urethral mucosal lining, Urethral hypermobility, Detrusor overactivity, Known polyuria, Unevaluated haematuria.

Side Effects: The following side effects may be associated with the use of the device system: General side effects normally associated with any surgical implantation procedure or local anaesthesia. Postoperatively, transient symptoms such as dysuria, stranguria, haematuria, urinary tract infection, and acute retention may occur; Scratching of the urethral mucosal may occur during the procedure; Long-term side effects such as non-acute retention, abscess formation, fibrosis (tissue hardening), de novo urgency, and necrosis are possible, but rare. Results and experiences may vary and are unique to each patient. No promise or guarantee is made about specific results or experiences. Talk to your doctor to find out if Bulkamid is right for you and to discuss the potential risks and benefits.

For a complete listing of indications, contraindications, warnings and precautions, go to www.bulkamid.com/isi
© 2021 Axonics, Inc. This material contains registered trademarks, trade names, and brand names of Axonics, Inc. All rights reserved. 110-0256-013rA 10/2021



Emergo Europe B.V. Prinsessegracht 20
2514 AP The Hague The Netherlands



Stay Connected.
#Axonics | #Bulkamid



What is stress urinary incontinence (SUI)?

Stress urinary incontinence (SUI), sometimes called stress incontinence, is a condition where unwanted bladder leaks happen during normal daily activities such as coughing, laughing, or exercise.

How can Bulkamid help?

Bulkamid is a urethral bulking agent that is used to treat stress urinary incontinence. It is a soft, water-based gel that can be used to restore the natural closing of the urethra. Similar to some facial fillers it remains in the body over time without causing reactions in the surrounding tissue.¹



Discover the benefits of Bulkamid

Safe and Effective

Bulkamid is a safe and effective treatment, with **92% of women** reporting being cured or improved.²

Simple Procedure

The procedure is a series of 3-4 small injections that takes around 10 to 15 minutes under local anaesthetic.³

Long-lasting Relief

Bulkamid is clinically proven to deliver symptom relief out to seven years.⁴

Talk to your doctor about Bulkamid today

The **majority of women** with SUI choose Bulkamid before other treatment options⁵

“ ”

I didn't experience any pain or discomfort and the effects of the treatment were almost immediate. No more leaks! I won't hesitate to recommend Bulkamid and the difference it's made to my life.

— Jodie M. | Bulkamid Patient, 42 years old —

“ ”

I haven't experienced another leak and I'd go as far to say it's changed my life. Now, I'm really spreading the word about my positive experience amongst those of my friends who might otherwise have carried on "suffering in silence."

— Natalie P. | Bulkamid Patient, 40 years old —

